

Lost at Sea

STEVEN R. FINCH AND JOHN A. SHONDER

November 23, 2004

ABSTRACT. What is the path of minimum expected length for escaping a planar convex region Ω ? We rigorously obtain best 2-segment and 3-segment solutions when Ω is an infinite strip, and numerically examine 2-segment solutions when Ω is a disk.

A swimmer is lost in a dense fog at sea. She knows that the sea is a planar infinite strip

$$\{(x, y) \in \mathbb{R}^2 : 0 \leq x \leq 1\}$$

of unit width. Assume that the x -coordinate of her initial position is uniformly distributed on the interval $[0, 1]$; the distribution of the y -coordinate can be arbitrary. Assume as well that her initial orientation (the angle between her initial velocity vector and the x -axis) is uniformly distributed on $[-\pi, \pi]$ and that her speed is constant. What escape trajectory should the swimmer follow that minimizes her expected time to reach either shore?

The min-max analog of this problem was solved long ago (see [1] for a survey). Zalgaller [2, 3] proposed a heuristic solution of the above min-mean problem. Shonder [4] independently determined the precise 2-segment solution of least expected escape time. (We agree that a k -segment path is a continuous, piecewise linear curve consisting of $\leq k$ pieces.) The details underlying Shonder's computation appear here for the first time. We find a 3-segment path that improves slightly upon the 2-segment path and also demonstrate that Zalgaller's solution is far from optimal.

Next, we examine a different sea: the disk

$$\{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\}$$

of unit radius. Assume that the starting point is uniformly distributed on the disk and that we seek (as before) to minimize the expected time to reach the boundary. Intuition suggests that the swimmer should follow a 1-segment path. If this is true, then the mean length of an arbitrary escape trajectory is $\geq 8/(3\pi) = 0.8488263631\dots$ [5, 6]. A calculus-of-variations proof of this general inequality is not known. We can confirm this only numerically for 2-segment paths.

In the final section, there appears a first attempt at proving the above inequality under special circumstances. The technique is due to Gevirtz [8] and yields results when the escape path curvature is sufficiently small.

⁰Copyright © 2004 by Steven R. Finch. All rights reserved.



Figure 1: Five cases for the infinite strip.

0.1. Infinite Strip: 2-Segment Scenario. Without loss of generality, let the initial position be $(x, 0)$ and the initial orientation be θ , where $0 \leq x \leq 1$ and $0 \leq \theta \leq \pi$. Fix a distance $r \geq 0$ and an angle $0 \leq \alpha \leq \pi$. As shown in Figure 1, there are five distinct cases to consider.

Case 1: the swimmer reaches the right-hand shore before traveling a distance r , which happens when $x + r \geq 1$ and $\theta \leq \arccos((1 - x)/r)$. The path length is $(1 - x)/\cos(\theta)$.

Case 2: the swimmer travels a distance r , does not reach the right-hand shore, pivots an angle α , and then travels until the left-hand shore is reached. This happens when

$$\{[x + r \geq 1 \text{ and } \arccos((1 - x)/r) < \theta] \text{ or } [x + r < 1]\} \text{ and } \{\theta < \frac{\pi}{2} - \alpha\}.$$

The path length is $r + (x + r \cos(\theta)) / \cos(\theta + \alpha)$.

Case 3: the swimmer travels a distance r , does not reach a shore, pivots an angle α , and then travels until the right-hand shore is reached. This happens when

$$\begin{aligned} & \{[x + r \geq 1 \text{ and } x - r \leq 0 \text{ and } \arccos((1 - x)/r) < \theta < \pi - \arccos(x/r)] \\ & \text{or } [x + r < 1 \text{ and } x - r \leq 0 \text{ and } \theta < \pi - \arccos(x/r)] \\ & \text{or } [x + r \geq 1 \text{ and } x - r > 0 \text{ and } \arccos((1 - x)/r) < \theta] \\ & \text{or } [x + r < 1 \text{ and } x - r > 0]\} \text{ and } \{\frac{\pi}{2} - \alpha \leq \theta \leq \frac{3\pi}{2} - \alpha\}. \end{aligned}$$

The path length is $r + (x - 1 + r \cos(\theta)) / \cos(\theta + \alpha)$.

Case 4: the swimmer travels a distance r , does not reach the left-hand shore, pivots an angle α , and then travels until the left-hand shore is reached. This happens when

$$\{[x - r \leq 0 \text{ and } \theta < \pi - \arccos(x/r)] \text{ or } [x - r > 0]\} \text{ and } \left\{\frac{3\pi}{2} - \alpha < \theta\right\}.$$

The path length is $r + (x + r \cos(\theta)) / \cos(\theta + \alpha)$.

Case 5: the swimmer reaches the left-hand shore before traveling a distance r , which happens when $x - r \leq 0$ and $\pi - \arccos(x/r) \leq \theta$. The path length is $-x / \cos(\theta)$.

We've restricted attention to $0 \leq \theta \leq \pi$ for simplicity's sake. The set of escape paths corresponding to $-\pi \leq \theta \leq 0$ is obtained from the set of escape paths corresponding to $0 \leq \theta \leq \pi$ via reflection across the horizontal axis, followed by reflection across the vertical axis. This composite transformation can be written as $(x, \theta) \mapsto (1 - x, \theta - \pi)$.

If it is assumed that $r > 1$ and $0 < \frac{\pi}{2} - \alpha < \arccos(1/r)$, then only Cases 1, 3 and 5 enter into the calculations of the expected path length:

$$\frac{1}{\pi} \int_0^1 \left[\int_0^{\arccos(\frac{1-x}{r})} \frac{1-x}{\cos(\theta)} d\theta + \int_{\arccos(\frac{1-x}{r})}^{\pi - \arccos(\frac{x}{r})} \left(r + \frac{x - 1 + r \cos(\theta)}{\cos(\theta + \alpha)} \right) d\theta - \int_{\pi - \arccos(\frac{x}{r})}^{\pi} \frac{x}{\cos(\theta)} d\theta \right] dx.$$

All of these integrals can be evaluated in closed form. Hence the expected path length, multiplied by π , becomes

$$\begin{aligned} & r \left[(1 + \cos \alpha) \left(\frac{\pi}{2} + \sqrt{r^2 - 1} - r \right) - (2 + \cos \alpha) \arccos \left(\frac{1}{r} \right) + \frac{\pi}{2} \right] + \\ & \ln \left(\sqrt{r^2 - 1} + r \right) + \frac{1}{2} r^2 \ln(1 - \cos \alpha) \sin^2 \alpha + r \sin \alpha \ln \left(\frac{\sqrt{r^2 - 1} \sin \alpha - \cos \alpha}{r \sin \alpha} \right) + \\ & \frac{1}{4} (r^2 \sin^2 \alpha + 1) \ln \left(\sqrt{r^2 - 1} \cos \alpha + \sin \alpha + r \right) + \\ & \frac{1}{4} (r^2 \sin^2 \alpha - 1) \ln \left(\sqrt{r^2 - 1} \cos \alpha - \sin \alpha + r \right) - \\ & \frac{1}{4} (r^2 \sin^2 \alpha + 1) \ln \left[-\sqrt{r^2 - 1} \cos^2 \alpha - \left(\sin \alpha + \sqrt{r^2 - 1} - r \right) \cos \alpha - \sin \alpha + r \right] - \\ & \frac{1}{4} (r^2 \sin^2 \alpha - 1) \ln \left[-\sqrt{r^2 - 1} \cos^2 \alpha + \left(\sin \alpha - \sqrt{r^2 - 1} + r \right) \cos \alpha + \sin \alpha + r \right]. \end{aligned}$$

Minimizing this expression gives $r = 1.0432668686\dots$ and $\alpha = 1.3734935859\dots \approx 78.7^\circ$ as the optimal parameter values for the 2-segment scenario. Therefore the least




Figure 2: Four sample realizations with $r = 1.043$, $\alpha = 78.7^\circ$, $x = 0.1, 0.3, 0.5, 0.9$ and $\theta = 80^\circ, 160^\circ, 50^\circ, 140^\circ$ (involving three Cases).




Figure 3: Two subcases of Case 3.

expected path length is 0.8869669056.... See Figure 2 for several sample realizations (with fixed r and α).

Here is a new problem. For each (r, α) , there exists a probability distribution of escape times obtained by sampling (x, θ) uniformly from $[0, 1] \times [0, \pi]$. Compute the median of each distribution and then determine (r, α) corresponding to the least median. Monte Carlo simulation suggests that a solution is $r = \infty$ (α can be arbitrary), consistent with the idea that medians are less sensitive to outliers than means. Define a random variable q to be $(1 - x)/\cos(\theta)$ if $\theta < \pi/2$ and $-x/\cos(\theta)$ if $\theta > \pi/2$. The path length is the median of q , which is estimated to be 0.78....

0.2. Infinite Strip: 3-Segment Scenario. We assume that $r > 1$ and $0 < \frac{\pi}{2} - \alpha < \arccos(1/r)$ as at the end of the preceding section. Cases 1 and 5 thus remain unchanged and only Case 3 needs to be refined. Fix a second distance $s \geq 0$ and a second angle $0 \leq \beta \leq \pi$. See Figure 3.

Subcase 3.1: the swimmer travels a distance r , does not reach a shore, pivots an angle α , and then reaches the right-hand shore before traveling a distance s . This happens when

$$\arccos((1 - x)/r) < \theta \leq \theta_0(x, r, s, \alpha)$$

where θ_0 is a solution of the equation

$$1 - x - r \cos(\theta) + s \cos(\theta + \alpha) = 0;$$

an exact formula for θ_0 is

$$\theta_0 = \arcsin \left(\frac{s \sin(\alpha)}{\sqrt{r^2 + s^2 - 2 r s \cos(\alpha)}} \right) + \arccos \left(\frac{1 - x}{\sqrt{r^2 + s^2 - 2 r s \cos(\alpha)}} \right).$$

The path length is $r + (x - 1 + r \cos(\theta)) / \cos(\theta + \alpha)$.

Subcase 3.2: the swimmer travels a distance r , does not reach a shore, pivots an angle α , travels a distance s , does not reach a shore, pivots an angle β , and then travels until the right-hand shore is reached. This happens when

$$\theta_0(x, r, s, \alpha) < \theta < \pi - \arccos(x/r)$$

and β satisfies other constraints (for example, if β is too small, then the 3-segment might cross itself and reach the left-hand shore rather than the right-hand shore). We will not attempt to prescribe these additional inequalities. The path length is $r + s + (1 - x + s \cos(\theta + \alpha) - r \cos(\theta)) / \cos(\theta + \alpha + \beta)$.

Under the proper circumstances, therefore, the expected path length is

$$\begin{aligned} \frac{1}{\pi} \int_0^1 \left[\int_0^{\arccos(\frac{1-x}{r})} \frac{1-x}{\cos(\theta)} d\theta + \int_{\arccos(\frac{1-x}{r})}^{\theta_0} \left(r + \frac{x-1+r \cos(\theta)}{\cos(\theta+\alpha)} \right) d\theta + \right. \\ \left. \int_{\theta_0}^{\pi - \arccos(\frac{x}{r})} \left(r + s + \frac{1-x+s \cos(\theta+\alpha)-r \cos(\theta)}{\cos(\theta+\alpha+\beta)} \right) d\theta - \int_{\pi - \arccos(\frac{x}{r})}^{\pi} \frac{x}{\cos(\theta)} d\theta \right] dx. \end{aligned}$$

A closed-form expression is unlikely here because of the presence of θ_0 in the middle two integrals: only integration with respect to θ seems feasible. Let

$$\kappa = \sqrt{r^2 + s^2 - 2 r s \cos(\alpha)}, \quad \rho = \arcsin(s \sin(\alpha)/\kappa),$$

$$u_1 = r \sin(\alpha), \quad u_2 = s \sin(\beta) - r \sin(\alpha + \beta),$$

$$v_1 = r(1 + \cos(\alpha)), \quad v_2 = r + s + s \cos(\beta) - r \cos(\alpha + \beta),$$

$$\xi_1 = \alpha + \arccos((1-x)/r), \quad \xi_2 = \alpha + \beta + \rho + \arccos((1-x)/\kappa),$$

$$\eta_1 = \alpha + \rho + \arccos((1-x)/\kappa), \quad \eta_2 = \alpha + \beta + \pi - \arccos(x/r)$$

and, for $j = 1, 2$, define

$$p_j = u_j \ln \left| \frac{\cos(\xi_j)}{\cos(\eta_j)} \right| + \frac{(-1)^j(1-x)}{2} \ln \left(\frac{\sin(\xi_j)(\sin(\eta_j) + 1) - \sin(\eta_j) - 1}{\sin(\xi_j)(\sin(\eta_j) - 1) + \sin(\eta_j) - 1} \right) + v_j(\eta_j - \xi_j).$$

Then the expected path length, multiplied by π , becomes

$$\ln(\sqrt{r^2 - 1} + r) + r(r - \sqrt{r^2 - 1}) + \int_0^1 (p_1 + p_2) dx.$$

This can be minimized by use of Leibniz's rule for differentiation under the integral sign. We obtain $r = 1.0255050653\dots$, $\alpha = 1.4909825316\dots \approx 85.4^\circ$, $s = 0.5306340577\dots$ and $\beta = 2.7495709960\dots \approx 157.5^\circ$ as the optimal parameter values for the 3-segment scenario, with least expected path length 0.8835534788....

A third subcase, for which $\pi \leq \beta \leq 2\pi$, awaits examination. Either shore could be the final resting place.

0.3. Infinite Strip: Zalgaller's Proposed Solution. Consider the path in Figure 4, where the points $A = (0, 0)$, $B = (0.814, 0)$, $C = (0.8460, 0.0005)$, $D = (1.3017, 0.0151)$, $E = (0.814, 1)$ are joined by line segments, with the exception of points B, C which are joined by a tiny circular arc of radius 1, center E and angle 0.032. Zalgaller [2, 3] claimed that, if the swimmer follows this path, then her expected time to reach either shore is approximately minimal. We examined his claim by computing best 2-segment and 3-segment fits to Zalgaller's path, and then calculating the expected escape time via our formulas. For example, in the 2-segment fit, we obtained $r = 1.3017$, $\alpha = 64.3^\circ$ and hence the expected escape time is 0.9188. This is somewhat consistent with Zalgaller's estimate 0.9523. More importantly, however, Shonder's optimal 2-segment path [4] has expected escape time 0.8870 which is considerably less than 0.9523. There must be an error somewhere in the details of [2]. We aim someday to better understand Zalgaller's elaborate construction, in the hope that his procedure (once corrected) will lead to escape trajectories that outperform even our optimal 3-segment path.

0.4. Disk: 2-Segment Scenario. Without loss of generality, let the starting point be $(x, 0)$ and the initial orientation be θ , where $0 \leq x \leq 1$ and $-\pi \leq \theta \leq \pi$. Fix a distance $0 \leq r \leq 2$ and an angle $0 \leq \alpha \leq \pi$. As shown in Figure 5, there are two distinct cases to consider. For convenience, define

$$\varphi(x, r) = \arccos\left(\frac{1 - x^2 - r^2}{2xr}\right), \quad \psi(x, r) = \arccos\left(-\frac{r}{x}\right).$$




Figure 4: Zalgaller's escape path.




Figure 5: Two cases for the disk.

Case 1': the swimmer reaches the shore before traveling a distance r , which happens when $x \geq |r - 1|$ and $-\varphi \leq \theta \leq \varphi$, or when $x < r - 1$. The path length is $q = -x \cos(\theta) + \sqrt{1 - x^2 \sin^2(\theta)}$.

Case 2': the swimmer travels a distance r , does not reach the shore, pivots an angle α , and then travels until the shore is reached. The path length is $r + s$, where

$$s = y \cos(\alpha + \omega) + \sqrt{1 - y^2 \sin^2(\alpha + \omega)},$$

$$y = \sqrt{x^2 + r^2 + 2 x r \cos(\theta)},$$

$$\omega = \begin{cases} \arcsin\left(\frac{x \sin(\theta)}{y}\right) & \text{if } (x \geq r \text{ and } -\psi \leq \theta \leq \psi) \text{ or } x < r, \\ \pi - \arcsin\left(\frac{x \sin(\theta)}{y}\right) & \text{if } x \geq r \text{ and } (\theta \leq -\psi \text{ or } \psi \leq \theta). \end{cases}$$

The proof of Case 1' makes use of the Law of Cosines and the fact that φ exists if and only if $x \geq |r - 1|$. The Law of Sines appears in the proof of Case 2' when defining the angle ω of an auxiliary triangle with side lengths r , x and y ; the Law of Cosines is then applied to the triangle with side lengths 1, s and y to obtain s .

The expected path length is therefore $I + J$, where

$$I = \frac{1}{\pi} \int_{|r-1|}^1 f(x, r, \alpha) x dx, \quad J = \begin{cases} \frac{1}{\pi} \int_0^{r-1} g(x) x dx & \text{if } r \geq 1, \\ \frac{1}{\pi} \int_0^{-r+1} h(x, r, \alpha) x dx & \text{if } r < 1 \end{cases}$$

and

$$f = \int_{-\varphi}^{\varphi} q d\theta + \int_{\varphi}^{\pi} (r + s) d\theta + \int_{-\pi}^{-\varphi} (r + s) d\theta, \quad g = \int_{-\pi}^{\pi} q d\theta, \quad h = \int_{-\pi}^{\pi} (r + s) d\theta.$$

Closed-form expressions are not possible here. Numerical computations confirm that $8/(3\pi)$ is the value of $I + J$ when $\alpha = \pi$, for any $r \geq 0$, and that otherwise $I + J$ is strictly larger than $8/(3\pi)$. See Figure 6 for several sample realizations (with fixed r but variable α).

We mention a corresponding min-median result: Monte Carlo simulation suggests that a solution is $r = 2$ (α can be arbitrary). The path length is the median of the random variable $-x \cos(\theta) + \sqrt{1 - x^2 \sin^2(\theta)}$, which is estimated to be 0.94....

These kinds of "lost at sea" problems deserve to be more thoroughly studied! The only other published reference we know regarding the min-mean problem is [7].




Figure 6: Sample realizations with $r = 0.5$, $x = 0.1, 0.3, 0.5$, $\alpha = 110^\circ, 160^\circ, 20^\circ$ and $\theta = 200^\circ, 80^\circ, 45^\circ$ (all from Case 2').

0.5. Disk: Gevirtz's Technique. Let $\gamma : [0, \infty) \rightarrow \mathbb{C}$ be a differentiable curve in the complex plane satisfying $\gamma(0) = 0$. Parametrize γ by arclength s and let $\gamma'(s) = \exp(i\phi(s))$, where we assume that $\phi(s) = \arg(\gamma'(s))$ is real, $\phi(0) = 0$ and $|\phi(s)| \leq \phi_0 < \frac{\pi}{2}$. The function $s \mapsto \lambda(s) = |\gamma(s)|$ is one-to-one and hence possesses an inverse $\lambda \mapsto s(\lambda)$. Let D denote the unit disk in $\mathbb{C}$ and define $C(z)$ to be the circle of radius 1 centered at z ; as an example, $C(0)$ is the boundary of D .

We wish to minimize the average

$$A(\gamma) = \frac{1}{\pi} \int_D \sigma(z, \gamma) dx dy$$

where $\sigma(z, \gamma)$ denotes the first (and only) value of s for which $|\gamma(s) - z| = 1$. (By symmetry of D , we have substituted $-z$ for z .) Now,

$$\int_D \sigma(z, \gamma) dx dy = \int_0^\infty s d\mu(s)$$

where $\mu(s)$ is the area of the portion of D not enclosed by $C(\gamma(s))$. This identity follows via a Riemann sum argument.

Consider the family of circular arcs $F = \{D \cap C(\gamma(s)) : 0 < s < \infty\}$. If ϕ_0 is small enough, then no two distinct arcs in F can intersect. As a consequence of the above identity, if the curvature $\phi'(s)$ is suitably small, then

$$A(\gamma) = \frac{2}{\pi} \int_0^{s^*} s \sqrt{1 - \left(\frac{|\gamma(s)|}{2}\right)^2} \cos(\phi(s) - \arg(\gamma(s))) ds$$

where s^* satisfies $|\gamma(s^*)| = 2$. Note that s^* depends on γ , which complicates any variational approach to this problem. Because $\lambda'(s) = \cos(\phi(s) - \arg(\gamma(s)))$ and $s(\lambda) \geq \lambda$, we have

$$A(\gamma) = \frac{2}{\pi} \int_0^2 s(\lambda) \sqrt{1 - \left(\frac{\lambda}{2}\right)^2} d\lambda \geq \frac{2}{\pi} \int_0^2 \lambda \sqrt{1 - \frac{\lambda^2}{4}} d\lambda = \frac{8}{3\pi},$$

as was to be shown. It is unclear what to do in the general situation for which distinct arcs in F are not necessarily disjoint. We also wonder if a similar technique exists for the infinite strip sea.

0.6. Acknowledgements. We are grateful to Julian Gevirtz and Patrice Le Conte for their help.

REFERENCES

- [1] S. R. Finch and J. E. Wetzel, Lost in a forest, *Amer. Math. Monthly* 11 (2004) 645-654.
- [2] V. A. Zalgaller, A discussion about a question of Bellman (in Russian), *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov (POMI)* 299 (2003) 54-86, 327-328; Engl. transl. available from S. Finch upon request (*J. Math. Sci.*, to appear?); MR2038255 (2004i:90174).
- [3] V. A. Zalgaller, Extremal problems concerning the convex hull of a space curve (in Russian), *Algebra i Analiz*, v. 8 (1996) n. 3, 1-13; Engl. transl. in *St. Petersburg Math. J.* 8 (1997) 369-379; MR1402285 (97c:58028).
- [4] J. A. Shonder, challenge and solution statement in F. Morgan, Math Chat, http://www.maa.org/features/mathchat/mathchat_8_3_00.html.
- [5] R. Coleman, Random paths through convex bodies, *J. Appl. Probab.* 6 (1969) 430-441; MR0254890 (40 #8097).
- [6] P. R. Coulton, Expectation relations for escape problems, unpublished note (2003).

- [7] B. Gluss, An alternative solution to the “lost at sea” problem, *Naval Res. Logist. Quart.* 8 (1961) 117–121; MR0129005 (23 #B2042).
- [8] J. Gevirtz, private communication (1999).

Steven R. Finch
Clay Mathematics Institute
One Bow Street
Cambridge, MA 02138
Steven.Finch@inria.fr

John A. Shonder
Oak Ridge National Laboratory
P.O. Box 2008, Bldg 3147, MS-6070
Oak Ridge, TN 37831
shonderja@ornl.gov

This figure "fig1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/math/0411518v1>

This figure "fig2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/math/0411518v1>

This figure "fig3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/math/0411518v1>

This figure "fig4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/math/0411518v1>

This figure "fig5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/math/0411518v1>

This figure "fig6.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/math/0411518v1>